

## Ultrafast Excited-State Dynamics in the Green Fluorescent Protein Variant S65T/H148D. 1. Mutagenesis and Structural Studies<sup>†,‡</sup>

Xiaokun Shu,<sup>§</sup> Karen Kallio,<sup>§</sup> Xinghua Shi,<sup>#</sup> Paul Abbyad,<sup>#</sup> Pakorn Kanchanawong,<sup>#</sup> William Childs,<sup>#</sup> Steven G. Boxer,<sup>#</sup> and S. James Remington<sup>\*,§</sup>

*Institute of Molecular Biology and Department of Physics, University of Oregon, Eugene, Oregon 97403-1229, and Department of Chemistry, Stanford University, Stanford, California 94305-5080*

*Received May 11, 2007; Revised Manuscript Received August 16, 2007*

**ABSTRACT:** Wild type green fluorescent protein (wt-GFP) and the variant S65T/H148D each exhibit two absorption bands, A and B, which are associated with the protonated and deprotonated chromophores, respectively. Excitation of either band leads to green emission. In wt-GFP, excitation of band A (~395 nm) leads to green emission with a rise time of 10–15 ps, due to excited-state proton transfer (ESPT) from the chromophore hydroxyl group to an acceptor. This process produces an anionic excited-state intermediate I\* that subsequently emits a green photon. In the variant S65T/H148D, the A band absorbance maximum is red-shifted to ~415 nm, and as detailed in the accompanying papers, when the A band is excited, green fluorescence appears with a rise time shorter than the instrument time resolution (~170 fs). On the basis of the steady-state spectroscopy and high-resolution crystal structures of several variants described herein, it is proposed that in S65T/H148D, the red shift of absorption band A and the ultrafast appearance of green fluorescence upon excitation of band A are due to a very short ( $\leq 2.4$  Å), and possibly low-barrier, hydrogen bond between the chromophore hydroxyl and introduced Asp148.

Wild type green fluorescent protein (wt-GFP)<sup>1</sup> exhibits two bands in absorption spectra, peaked at 395 and 475 nm and designated A and B, respectively [see reviews (3–5)]. Whereas band A is attributed to absorption of the protonated chromophore, band B is due to absorption of the deprotonated chromophore (6, 7). Excitation of either band leads to green emission; however, upon excitation of band A, the intensity of green emission rises on a time scale of several tens of picoseconds. This process is dramatically slowed upon deuteration of the sample, suggesting that excited-state proton transfer (ESPT) from the neutral form of the chromophore generates an intermediate anionic excited state, which subsequently emits a green photon (6–9). On the basis of the crystal structures, a pathway has been proposed to allow proton transfer between the chromophore phenol moiety, through a water molecule and Ser205, to Glu222 as the terminal proton acceptor (10–12). In wt-GFP, the intensities

of absorbance bands A and B are largely insensitive to pH over the range 4–10.

Point mutations can lead to increased sensitivity to pH and the formation of new ESPT pathways within GFP. For example, dual emission GFPs (deGFPs) switch between a form that emits blue light at low pH and a form that emits green light at high pH. Time-resolved spectroscopy and structural studies of deGFPs suggested that the dual emission behavior results from pH-dependent formation of alternative ESPT pathways (13–15). Mutations can also eliminate the fluorescence from band A; for example, the S65T variant [a precursor of a popular molecular label EGFP (16)], has a strong pH dependence of fluorescence with  $pK_a$  of about 6.0. At pH <6.0 the protonated form of the chromophore predominates, leading to strong absorbance from band A; however, fluorescence emission from the protonated chromophore is vanishingly weak.

The subject of this and the accompanying papers (1, 2) is the unusual photophysical behavior of the double-mutant GFP S65T/H148D, the construction of which was previously described (17). As briefly described in that work, the additional mutation His148 → Asp upon the S65T background restores green fluorescence from band A (17). As with the variant S65T, the excitation and emission spectra of S65T/H148D are pH sensitive, consistent with a chromophore  $pK_a$  of about 8.0. As the pH is varied, two-state behavior is observed with respect to green emission from either band A or band B, indicating that the two species

\* Author to whom correspondence should be addressed [telephone (541) 346-5190; fax (541) 346-5870; e-mail jremington@uoxray.uoregon.edu].

<sup>†</sup> This work was supported by grants from the NIH (R01 GM42618 to S.J.R. and GM 27738 to S.G.B.).

<sup>‡</sup> The atomic coordinates and structure factors have been deposited in the Protein Data Bank (entry 2DUF and 2DUE for GFP S65T/H148D at pH 5.6 and 10.0 respectively, 2DUG and 2DUH for GFP S65T/H148N at pH 5.0 and 9.5, 2DUI for GFP H148D at pH 9.0).

<sup>§</sup> University of Oregon.

<sup>#</sup> Stanford University.

<sup>1</sup> Abbreviations: GFP, green fluorescent protein; H-bond, hydrogen bond; LBHB, low-barrier hydrogen bond; ESPT, excited-state proton transfer; IRF, instrument response function; rms, root-mean-square.

Table 1: Effects of Substitution of Residue 148 on the GFP

mutant	absorbance band (nm)		emission band <sup>a</sup> (nm)		pK <sub>a</sub>	quantum yield <sup>b</sup>	
	A	B	A	B		A	B
S65T	394	489	459/508	510	5.95 ± 0.02 <sup>c</sup>	0.02 ± 0.01	0.67 ± 0.01
S65T/H148N	393	488	459/510	511	6.37 ± 0.02	0.01 ± 0.01	0.45 ± 0.02
S65T/H148E	397	484	460/510	511	7.17 ± 0.05	0.04 ± 0.01	0.24 ± 0.02
S65T/H148D	415	487	510	510	7.95 ± 0.02 <sup>c</sup>	0.21 ± 0.04	0.32 ± 0.03
H148D	400	469	503	505	>10.4	0.15 ± 0.02	0.40 ± 0.02
wild type GFP	395	475	460/509	505	NA	0.75 ± 0.03	0.75 ± 0.03

<sup>a</sup> Emission maxima resulting from excitation at the peak of either absorbance band A or B. <sup>b</sup> Quantum yields of bands A and B were measured at pH 4.5 or 5 and 11, respectively. <sup>c</sup> Taken from ref 17.

directly interconvert. Thus, this mutant finds application as a ratiometric pH sensor (18).

GFP S65T/H148D exhibits unusual photophysical properties. Absorbance band A is maximal at 415 nm (17); thus, it is red-shifted by ~20 nm compared to wt-GFP or the S65T variant. In time-resolved spectroscopic studies, both weak blue (~460 nm) and strong green (~510 nm) emissions are observed upon excitation of band A at 400 nm; however, both blue and green emissions are characterized by a rise time that is shorter than the instrument response function (IRF, ~170 fs). As detailed in the accompanying papers (1, 2), the emission kinetics are only weakly sensitive to deuterium substitution.

Here, we report on mutagenic and high-resolution crystallographic studies which demonstrate that the unusual photophysical properties of the S65T/H148D variant are related to a very short hydrogen bond ( $\leq 2.4$  Å) between the chromophore hydroxyl and Asp148.

## MATERIALS AND METHODS

**Mutagenesis, Protein Expression, and Crystallization.** Mutagenesis was performed using the QuikChange method (Stratagene). Protein was expressed in *Escherichia coli* strain JM109 (DE3), by use of the PRSET<sub>B</sub> His tagged expression system. Protein was purified by Ni<sup>2+</sup> affinity chromatography over Ni-NTA agarose (Qiagen, Chatsworth, CA) and then buffer exchanged with PD-10 Sephadex columns (Amersham Pharmacia) into 50 mM HEPES (pH 7.9). Crystals of S65T/H148D were obtained after 2 months by hanging drop vapor diffusion [2 μL of protein solution (50 mM HEPES, pH 7.9, A<sub>280</sub> = 20.3) was mixed with 2 μL of well solution (100 mM MgCl<sub>2</sub>, 100 mM NaOAc, pH 5.6, 26% PEG 4000)]. For the high-pH structure of S65T/H148D, crystals were soaked in 100 mM MgCl<sub>2</sub>, 100 mM CAPS, pH 10.0, and 26% PEG 4000 overnight. S65T/H148N crystallized in 50 mM Li<sub>2</sub>SO<sub>4</sub>, 100 mM Tris, pH 9.5, and 30% PEG 4000 after 1 week. To obtain crystals of S65T/H148N at low pH, crystals were soaked in 200 mM Ca(OAc)<sub>2</sub>, 100 mM MES, pH 5.0, and 20% PEG 8000. GFP H148D crystallized in 100 mM MgCl<sub>2</sub>, 100 mM Tris, pH 9.0, and 30% PEG 1550 overnight.

**Spectroscopy.** Absorbance spectra (400 μg/mL protein in 100 mM buffers) were recorded with a HP 8453 UV–visible spectrophotometer. Protein pK<sub>a</sub> values were determined as described previously (13). Fluorescence spectra were taken with a Perkin-Elmer LS-55 fluorometer after dilution of protein samples to 30–100 μg/mL in 2 mL of 100 mM buffers of citric acid, HEPES, or CHES as appropriate. The quantum yields (QY) of emission were measured by refer-

encing to the standards 9-aminoacridine and fluorescein. As controls, QY were determined for wt-GFP and S65T as well (Table 1). These values are in good agreement with previous results (4, 5).

**Data Collection and Structure Solution.** Diffraction data sets were collected from single flash-frozen crystals of S65T/H148N at pH 5.0 and 9.5 (100 K) using ADSC-Q315 detector on beam line 8.2.2 at the Advanced Light Source (Berkeley, CA). Diffraction data for S65T/H148D at pH 5.6 and 10.0, H148D at pH 9.0, were collected on a Quantum-315 CCD on beam line 14-BM-C at the Advanced Photon Source (Argonne, IL). Data were reduced using HKL2000 (HKL Research) or d\*TREK (Molecular Structure Corp.). Molecular replacement was performed with EPMR (19) using the GFP S65T (PDB entry 1EMA) as a search model, providing unambiguous identification of the space group and unique solutions to the rotation and translation problems. Rigid body refinement was performed using TNT (20), and model building was conducted with O (21) or Coot (22) in several stages of increasing resolution. SHELX-97 (23) was used in the final stages of refinement for S65T/H148D at pH 10.0, S65T/H148N at pH 9.5, and H148D at pH 9.0. To calculate the standard uncertainties of the very short hydrogen bond between the chromophore and Asp148 (see Results), the low-pH S65T/H148D model was also submitted to automated refinement by SHELX.

## RESULTS

**Steady-State Spectra.** GFP S65T/H148D displays two pH titratable (pK<sub>a</sub> ~ 8.0) bands in absorption and excitation spectra (Figure 1a,b) (17). Compared to the S65T variant, band A is broad and red-shifted to 415 nm (by ~20 nm). However, the B band maximum remains at ~490 nm, essentially unchanged from that of S65T. Excitation of band A leads to relatively strong green emission (QY = 0.21), which is very weak in the S65T variant (QY ~ 0.02). Evidently, upon band A excitation, the H148D mutation partially reverses the severe reduction in fluorescence efficiency attributed to the S65T mutation. The emission maximum is approximately the same (509 ± 1 nm) for all variants except for GFP/H148D, which emits at ~503 nm.

To investigate the role of Asp148 on the spectroscopic properties we first mutated it to asparagine, which is isosteric but cannot carry a charge. The S65T/H148N variant has properties that are very similar to those of S65T (Table 1). Absorbance spectra reveal two pH-titratable bands (Figure 1a) with pK<sub>a</sub> ~ 6.4, suggesting that a partial charge on Asp148 could be responsible for the apparent chromophore pK<sub>a</sub> of 8.0 in S65T/H148D. The Asn148 substitution restores




FIGURE 1: Normalized absorbance, excitation, and emission spectra: (a) absorbance spectra of S65T/H148E/N/D at pH 7.3/6.1/7.6, respectively; (b) excitation spectra of S65T/H148E/N/D by monitoring fluorescence at 510 nm at pH 6.2/5.4/7.0, 1 unit lower than the respective variants'  $pK_a$  values; (c) excitation spectra of H148D with detection at 508 nm as a function of pH; (d) emission spectra of S65T/H148E with excitation at 400 nm as a function of pH.

the band A absorbance maximum to  $\sim 395$  nm, whereas the band B maximum remains at  $\sim 490$  nm. S65T/H148N is essentially nonfluorescent ( $QY \sim 0.01$ ) upon excitation of band A (Figure 1b). Excitation of band B, however, leads to efficient green fluorescence with  $QY = 0.45$ .

The possible effect of a negative charge at the 148 position was studied by substitution with glutamate. The variant S65T/H148E also shows two bands in absorption spectra (Figure 1a) with the band A maximum again consistent with wild type at  $\sim 395$  nm. Green emission from band A is weak with  $QY = 0.04$ , whereas emission from band B has  $QY = 0.24$  (Table 1). Nevertheless, S65T/H148E is a new dual-emission GFP [deGFP (13)], in that it shows both blue and green emissions upon excitation at 400 nm, with relative intensities depending strongly on pH (Figure 1d). Blue emission at 460 nm is increased at low pH, whereas the green emission is increased at high pH. The  $pK_a$  (7.2) of S65T/H148E is very close to the physiological pH and is intermediate between those of S65T/H148D (8.0) and S65T/H148N (6.4). Thus, this mutant might find application as a pH indicator in living cells.

For these mutations, the change in side-chain polarity near the chromophore (specifically at positions 148 and/or 222) correlates well with an increase in the chromophore  $pK_a$ , as exhibited by the series S65T  $\rightarrow$  S65T/H148N  $\rightarrow$  S65T/H148E, S65T/H148D  $\rightarrow$  H148D (Table 1).

We also characterized the role of the S65T mutation in the S65T/H148D variant by reverting Thr65 to serine. The generated variant H148D behaves in many ways like wt-GFP. The A band maximum is 400 nm in both absorption and excitation spectra (for emission at 510 nm). Excitation of either the A or B band leads to green emission, as in wt-GFP (Figure 1c). Band A dominates at pH below 9 and can be decreased by increasing pH higher than 9, with apparent  $pK_a > 10.5$ . This variant is ratiometric by excitation (Figure 1c), similar to S65T/H148D (17). Thus, the mutation His148  $\rightarrow$  Asp on the wild-type background is shown to confer ratiometric pH-sensitive behavior, which might conceivably find applications in high-pH environments; however, the  $pK_a$  is very high.

We conclude from the mutagenesis and the steady-state spectroscopic analysis that in S65T/H148D, the H148D

Table 2: Data Collection and Refinement Statistics<sup>a</sup>

	S65T/H148D (pH 5.6)	S65T/H148D (pH 10.0)	S65T/H148N (pH 5.0)	S65T/H148N (pH 9.5)	H148D (pH 9.0)
data collection					
space group	<i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	<i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	<i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	<i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	<i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>
cell dimensions <i>a</i> , <i>b</i> , <i>c</i> (Å)	51.1, 63.0, 68.7	50.9, 62.6, 69.9	51.2, 62.5, 68.9	50.7, 62.4, 68.6	50.7, 62.6, 68.6
resolution (Å)	50–1.50 (1.55–1.50)	50–1.24 (1.28–1.24)	34–1.40 (1.45–1.40)	10–1.20 (1.24–1.20)	50–1.36 (1.41–1.36)
<i>R</i> <sub>sym</sub> or <i>R</i> <sub>merge</sub> (%)	4.9 (42.3)	4.4 (32.9)	5.5 (50.8)	5.3 (33.4)	4.9 (31.9)
<i>I</i> / <i>σI</i>	35.0 (2.1)	42.9 (2.7)	11.0 (2.0)	17.5 (3.9)	55.7 (8.4)
completeness (%)	94.7 (68.6)	93.7 (61.8)	98.1 (85.9)	94.5 (93.3)	96.0 (86.9)
redundancy	3.2 (2.2)	3.5 (2.2)	6.4 (3.9)	6.5 (6.2)	3.7 (3.6)
refinement					
resolution (Å)	6–1.50	10–1.24	6–1.40	10–1.20	10–1.36
no. of reflections	34,193	59,945	43,438	61,533	45,807
<i>R</i> <sub>work</sub> / <i>R</i> <sub>free</sub>	17.2/24.4	15.47/20.4	17.7/24.1	15.2/20.9	15.1/21.6
no. of atoms					
protein	1769	1773	1799	1808	1797
water	294	249	237	324	267
<i>B</i> factors (Å <sup>2</sup> )					
protein	27.0	18.1	28.7	19.0	21.9
water	40.1	32.0	43.8	37.4	36.5
rms deviations					
bond length (Å)	0.017	0.013	0.020	0.013	0.011
bond angle (Å)	1.8(°)	0.031	1.9(°)	0.031	0.029

<sup>a</sup> Values in parentheses indicate statistics for the highest resolution shell.

mutation is required both for the ~20 nm red shift of band A and for green emission from band A. The S65T mutation is required for the red shift, acting synergistically with H148D; however, S65T is not essential for green emission upon excitation of band A.

**Crystal Structures.** As detailed in the accompanying papers (1, 2), the photophysics of green emission from S65T/H148D (upon band A excitation) is unusual for its rise time of <170 fs (IRF). To investigate the structural basis for this remarkable behavior, crystal structures of S65T/H148D, S65T/H148N, and H148D were solved, with the former two at both low and high pH. All variants crystallized isomorphously with S65T GFP (24), with one protomer in the asymmetric unit. Data collection and refinement statistics for the atomic models are presented in Table 2. In all crystals, electron density is weak for residues 230–238, so these residues were not modeled. For the same reason, residues not modeled in S65T/H148D were 1–3, 189–190 (low pH), and 1, 147–148, 155–157 (high pH); in S65T/H148N those residues not modeled were 1–3, 156–157 (low pH), and 1 (high pH); and those residues not modeled in H148D were 1–3, 147–148, and 155–159 (pH 9.0). The refinement and atomic model statistics are excellent. PROCHECK (25) reveals that there are no residues in either the disallowed or “generously allowed” regions of the Ramachandran diagram.

Upon superposition of the  $\alpha$ -carbons with GFP S65T, root-mean-square (rms) deviations are observed to be <0.40 Å for all structures. The atomic models are thus all very similar, which indicates that the protein matrix is largely insensitive toward variation at position 148. Schematic diagrams (Figure 2a–c) of the chromophore environment in each case demonstrate that the overall environment is similar to that of S65T or wt-GFP. Nevertheless, interesting and significant local effects are observed as detailed in the following sections.

(a) *S65T/H148D*. The crystal structure of this protein at pH 5.6 was refined at 1.5 Å resolution. A portion of the final 2Fo-Fc map (Figure 3) reveals that a very short hydrogen bond, 2.32 Å (as determined after refinement using

isotropic temperature factors with TNT), forms between the phenol oxygen of the chromophore and a carboxylate oxygen of Asp148. After SHELX refinement with anisotropic temperature factors, the length of this hydrogen bond is estimated to be  $2.41 \pm 0.03$  Å (estimated standard error). Given that the model errors in the two refinement procedures are different, the most probable value lies somewhere between these two estimates. Asp148 adopts the least favored  $\chi_1$  rotamer (~60°, see Figure 3), which, according to the rotamer library of COOT (22), is seen in only 2% of cases. Two symmetrically placed water molecules are hydrogen bonded with the two carboxylate oxygens of Asp148 separately. In addition, a hydrogen-bonded water chain is attached to Asp148, leading to the exterior of the protein. Finally, a hydrogen bond network (chromophore OH  $\rightarrow$  H<sub>2</sub>O  $\rightarrow$  Ser205  $\rightarrow$  Glu222) is formed, restoring the potential ESPT pathway found in wt-GFP (10, 11). As this ESPT pathway is disrupted by the S65T mutation, it appears that one consequence of the H148D mutation is indeed to reverse some of the changes caused by the S65T mutation, even though these sites are spatially quite distinct in the molecule.

The crystal structure of S65T/H148D at pH 10 was determined at 1.24 Å resolution. A dramatic structural change is observed in that the main-chain segment comprising Asp148 and Ser147 is disordered at this pH. No significant electron density is observed in the final 2Fo-Fc map for these two residues (data not shown). We presume that the disorder is a result of a repulsive electrostatic interaction between the negatively charged chromophore and Asp148. As described below, this observation accounts for the overall pH dependence of fluorescence.

(b) *S65T/H148N*. Crystal structures of this protein were determined at 1.4 (pH 5.0) and 1.2 Å resolution (pH 9.5). The low-pH crystal structure indicates that Glu222 is partially disordered and has two conformations, one of which is found in the S65T structure (Figure 2b) (17), whereas the other conformation restores a potential ESPT pathway similar to that found in wt-GFP (chromophore OH  $\rightarrow$  H<sub>2</sub>O  $\rightarrow$  Ser205  $\rightarrow$  Glu222) (10, 11). However, this pathway seems not to








FIGURE 4: Stereo image of superposed atomic models showing structural changes in S65T/H148N at pH 5.0 (black bonds) and 9.5 (green bonds). Dashed lines indicate hydrogen bonds, labeled with distances in angstroms. Blue spheres are nitrogen; red are oxygen.




FIGURE 5: Stereo image of (2Fo-Fc) electron density map and atomic model of H148D at pH 9.0 showing potential ESPT pathway; contour level  $1\sigma$ . Blue spheres are nitrogen; red are oxygen. Dashed lines indicate the hydrogen bonds, labeled with distances in angstroms. For other hydrogen bond lengths refer to Figure 2c.

Table 3: Chromophore Geometry in the GFPs, Angles in Degrees<sup>a</sup>

model	QY	twist	tilt
wt-GFP	0.79	0.0	3.7
DsRed	0.79	0.9	0.2
S65T/H148D (pH 5.6)	0.21	9.5	9.6
S65T/H148D (pH 10.0)	0.32	3.2	9.0
S65T/H148N (pH 5.0)	0.01	7.6	13.5
S65T/H148N (pH 9.5)	0.45	1.3	7.7
H148D (pH 9.0)	0.15	0.4	12.8

<sup>a</sup> See ref 27 for a definition of twist and tilt angles. Quantum yields (QY) were measured by excitation at maximum of absorbance band at specific pH.

tilt angles describing the distortion are significantly larger than those of the highly fluorescent proteins such as DsRed and GFP (27) (Table 3), but less than those found in the nonfluorescent chromoproteins KFP (27) and Rtm5 (28). The chromophore for both S65T/H148D and S65T/H148N is more bent at low pH than at high pH, which correlates with the dependence of the QY on pH (Table 1). This apparent correlation between relatively low QY and the chromophore non-coplanarity is also seen in monomeric red fluorescent proteins known as mFruits (29), suggesting a general relationship. The non-coplanarity of the chromophore is presumably due to the shape of the protein cavity (30).

## DISCUSSION

The double-mutant GFP S65T/H148D is unusual with respect to a number of its photo- and biophysical properties. The introduction of the S65T mutation into wt-GFP leads to loss of fluorescence upon excitation of the A band, which is restored by the second site mutation H148D. Unlike wt-GFP, both mutants are strongly pH sensitive and display two-state titration behavior. S65T/H148D is ratiometric by excitation, which potentially makes this variant useful as a pH sensor for use in living cells. Furthermore, the second

site mutation introduces a large red shift into the A band absorbance, without introducing a significant shift in the B band absorbance maximum or the fluorescence emission. Finally, the rise time of green emission is ultrafast, in contrast to wild type, for which green emission upon excitation of the A band shows a rise time of tens of picoseconds.

Structural studies revealed that in S65T/H148D, Asp148 makes a very short ( $<2.4$  Å) hydrogen bond with the chromophore hydroxyl oxygen, which suggests a strong, specific interaction. However, it is possible that the effect of Asp148 is due to a nonspecific electrostatic interaction; that is, the effect results from the introduction of a negative charge adjacent to the chromophore. To examine these possibilities and untangle the various possible contributions of Asp148 and Thr65, we made a number of substitutions at these positions and compared their respective biophysical properties.

In the following we focus first on the structural basis of the ratiometric pH sensitivity and then on the possible relationships between the very short H-bond and the unusual photophysical properties of the S65T/H148D mutant. Several possible mechanisms accounting for the extremely short rise time of green fluorescence are briefly discussed. For more details, the reader is referred to the two accompanying papers (1, 2).

*Structural Basis of the Ratiometric pH Sensitivity.* pH-dependent transitions in the excitation spectra of S65T/H148D can be accurately modeled by assuming a single-site titration with  $pK_a \sim 8.0$ . The structural basis of the ratiometric pH sensitivity is suggested by the crystal structures of S65T/H148D at pH 5.6 and 10.0. At low pH, a well-ordered H-bond forms between the chromophore and Asp148, whereas at high pH, the chain segment composed of Ser147 and Asp148 becomes highly disordered. It is expected that the  $pK_a$  of Asp148 is much lower than 8.0, and thus it is reasonable to assign the apparent  $pK_a$  of  $\sim 8.0$

to direct titration of the chromophore hydroxyl group. This assignment is supported by evidence from structural analysis of S65T/H148N at low and high pH, where the pH-dependent rearrangement of Asn148 and Thr203 implicates the chromophore hydroxyl as the titrating group, with  $pK_a$  of about 6.4. The assignment is also consistent with earlier structural studies of the GFP variant S65T, in which the chromophore hydroxyl was identified as the titrating group (17).

A simple two-state model adequately explains the ratio-metric response of S65T/H148D to titration. At low pH, the chromophore population is on average protonated, and as in wt-GFP, ESPT can explain the strong green fluorescence from band A. At high pH, the anionic form of the chromophore predominates and gives rise to green fluorescence as expected. Due to charge–charge repulsion between Asp148 and the chromophore, a segment of the main chain comprising residues 147 and 148 becomes disordered. This disorder does not appear to substantially compromise fluorescence efficiency for excitation of the anionic chromophore.

However, the results from ultrafast spectroscopy suggest that for wt-GFP and S65T/H148D, the mechanisms of fluorescence emission upon excitation of the A bands are different.

**Band A Red Shift.** As described above, the red shift of band A of GFP S65T/H148D can be attributed to the interaction between the chromophore and Asp148. To determine whether the red shift is due to an electrostatic interaction or is related to the formation of a short hydrogen bond, we introduced further mutations at positions 65 and 148 and characterized the resulting purified proteins.

Substitution of Asp148 with glutamate leads to normal A band absorbance of 397 nm, observable over the range of pH 5–9 (above pH 9, A band absorbance is weak or nonexistent). It is very likely that the introduced Glu148 is negatively charged over this range; thus, we conclude that a presumed negative charge on Asp148 cannot be responsible for the red shift in the S65T/H148D mutation. Likewise, reversion of Thr65 to Ser, as found in wild type, leads to a variant containing the single mutation H148D. This variant has a normal A band absorbance maximum at 400 nm and an extremely high apparent  $pK_a$  ( $>10$ ). This behavior is similar to that of wt-GFP, in which the negative charge of Glu222 is proposed to stabilize the protonated chromophore (10). Thus, it is likely that in H148D, Glu222 is also negatively charged. Structural studies revealed that the Asp148 side chain is disordered but is, nevertheless, in the immediate vicinity of the chromophore. It is very likely that Asp148 is anionic over the range of pH 9–11. Thus, the absence of a significant red shift is consistent with our previous conclusion that electrostatic effects alone cannot be responsible for the effect.

Finally, the mutant S65T/H148N was investigated, and crystal structures were determined at pH 5.0 and 9.5. Asn148 forms a hydrogen bond to the chromophore hydroxyl at either pH (Figure 4), but in either case the hydrogen bond lengths are typical and the A band absorbance maximum is about 393 nm. However, the side chain of Asn148 does rearrange as a function of pH, consistent with a change in hydrogen bonding due to dissociation of the chromophore hydroxyl at high pH.

From the above, we conclude that in the variant S65T/H148D, the band A red shift is due to the short hydrogen

bond formed between the chromophore and Asp148. One possible explanation for the red shift is that the short hydrogen bond promotes additional delocalization of the chromophore electron density over the carboxylate of Asp148, thus increasing the physical extent of the chromophore. If this interaction were to be maintained in the excited state, one would expect emission to be similarly red-shifted; however, it is normal at  $\sim 510$  nm. On the basis of this observation, we additionally suggest that the chromophore–Asp148 interaction is altered or lost in the excited state.

**Ultrafast Rise Time of Green Fluorescence.** Upon excitation of the A band, the rise time of green fluorescence of S65T/H148D is faster than the instrument response function (170 fs). Weak blue fluorescence is also observable; however, as detailed in the accompanying papers (1, 2), the kinetics of blue and green fluorescence suggest multiple emitting species, some of which may not interconvert via ESPT pathways. As pointed out above, in S65T/H148D an ESPT pathway similar to that found in wt-GFP is present (see Figure 3). It seems unlikely that the restored ESPT pathway could be responsible for the ultrafast kinetics of green emission of S65T/H148D, because in wt-GFP, green fluorescence attributed to ESPT arises on the time scale of tens of picoseconds (6, 7). Moreover, the rise time is significantly slowed upon deuteration of the sample, which is not the case for S65T/H148D.

Thus, one must look elsewhere to explain the ultrafast rise time ( $<170$  fs) of green fluorescence. We again suggest that the short H-bond between the chromophore and Asp148 is responsible for this behavior. One possibility to consider is ESPT from the protonated chromophore phenol to Asp148 through the short H-bond. Such ultrafast proton transfer (with transfer times expected to be on the order of 10–20 fs) has previously been described, for example, in solution between preformed, hydrogen-bonded HPTS $\cdots$ OAc complexes (31). However, to our knowledge, there is no precedence for similar observations of ultrafast proton transfer within a protein molecule.

A second possibility is that the short hydrogen bond indeed has a low barrier, or no barrier, to proton transfer and that the proton is delocalized between the chromophore hydroxyl and the carboxylate of aspartate 148. Thus, effectively instantaneous proton transfer could be achieved as a result of collapse of the proton wave function.

As detailed in the accompanying papers (1, 2), the kinetics of blue and green fluorescence in S65T/H148D are quite complicated. This may be related to the presence of a minor species or proton-transfer pathway not apparent in the crystal structure (up to 10–15% occupancy of a minor species would be difficult or impossible to detect in the electron density maps). It should be noted that the crystal contains no glycerol, whereas low-temperature spectroscopic experiments in the accompanying papers (1, 2) were conducted with the sample in  $\sim 60\%$  glycerol solution. As an example of alternative species not detectable in the crystal structure, different protonation or tautomeric states of the titratable groups comprising the chromophore–Asp148 interaction may need to be considered. The reader is referred to the accompanying papers (1, 2) for kinetic analyses and more detailed discussions.



*A Low-Barrier Hydrogen Bond?* To investigate whether the short hydrogen bond between the chromophore and Asp148 is a so-called low-barrier hydrogen bond (LBHB) (32, 33), we determined the 600 MHz jump-return  $^1\text{H}$  NMR spectrum of GFP S65T/H148D (pH 5.6, 28 °C). However, the spectrum does not reveal any unusually downfield-shifted resonances diagnostic of a solvent exchange-protected hydrogen in a LBHB (34, 35) (see Supporting Information Figure S1). The resolved signals between 10 and 13 ppm all arise from nitrogen-bonded protons in imidazole or amide moieties, as evidenced by  $\sim 95$  Hz scalar couplings in a uniformly  $^{15}\text{N}$ -labeled sample of the protein (L. McIntosh and M. Heller, data not shown). No additional downfield signals were detected upon reduction of the temperature to 5 °C. The excitation maximum for the jump-return pulse sequence was  $\sim 15$  ppm. We conclude from the NMR data that there is no evidence for an unusual downfield chemical shift related to any O–H $\cdots$ O hydrogen bond. It nevertheless seems to be possible that the very short hydrogen bond between the chromophore hydroxyl and Asp148 is a LBHB that is not protected from rapid solvent exchange. On the other hand, cases have been identified in which the chemical shift of a proton in a very short hydrogen bond is not as large as one might expect (36, 37). These observations were attributed to increased shielding of the proton by the approaching heavy atoms.

In conclusion, we suggest that the double-mutant GFP-S65T/H148D and several variations thereof offer unique opportunities to study excited-state proton-transfer pathways in an environment characterized by an unusually short hydrogen bond. Furthermore, the environment can be perturbed by mutagenesis, and the positions of the interacting groups may be determined at atomic resolution.

## ACKNOWLEDGMENT

We thank L. McIntosh and M. Heller, University of British Columbia, for obtaining the 1D-NMR spectra of S65T/H148D.

## SUPPORTING INFORMATION AVAILABLE

Figure S1, showing results from a 600 MHz jump-return NMR experiment to investigate whether unusual downfield shifts might be associated with the chromophore–Asp148 hydrogen bond. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## REFERENCES

- Shi, X., Abbyad, P., Shu, X., Kallio, K., Kanchanawong, P., Childs, W., Remington, S. J., and Boxer, S. G. (2007) Ultrafast excited-state dynamics in the green fluorescent protein variant S65T/H148D. 2. Unusual photophysical properties, *Biochemistry* 46, 12014–12025.
- Leiderman, P., Genosar, L., Huppert, D., Kallio, K., Remington, S. J., Solntsev, K. M., and Tolbert, L. M. (2007) Ultrafast excited-state dynamics in the green fluorescent protein variant S65T/H148D. 3. Short- and long-time dynamics of the excited-state proton transfer, *Biochemistry* 46, 12026–12036.
- Remington, S. J. (2000) Structural basis for understanding spectral variations in green fluorescent protein, *Bioluminescence Chemiluminescence, Part C* 305, 196–211.
- Tsien, R. Y. (1998) The green fluorescent protein, *Annu. Rev. Biochem.* 67, 509–544.
- Zimmer, M. (2002) Green fluorescent protein (GFP): Applications, structure, and related photophysical behavior, *Chem. Rev.* 102, 759–781.
- Chattoraj, M., King, B. A., Bublitz, G. U., and Boxer, S. G. (1996) Ultra-fast excited state dynamics in green fluorescent protein: multiple states and proton transfer, *Proc. Natl. Acad. Sci. U.S.A.* 93, 8362–8367.
- Lossau, H., Kummer, A., Heinecke, R., Pollinger-Dammer, F., Kompa, C., Bieser, G., Jonsson, T., Silva, C. M., Yang, M. M., Youvan, D. C., and Michel-Beyerle, M. E. (1996) Time-resolved spectroscopy of wild-type and mutant green fluorescent proteins reveals excited state deprotonation consistent with fluorophore–protein interactions, *Chem. Phys.* 213, 1–16.
- Creemers, T. M., Lock, A. J., Subramaniam, V., Jovin, T. M., and Volker, S. (1999) Three photoconvertible forms of green fluorescent protein identified by spectral hole-burning, *Nat. Struct. Biol.* 6, 557–560.
- Kennis, J. T. M., Larsen, D. S., van Stokkum, I. H. M., Vengris, M., van Thor, J. J., and van Grondelle, R. (2004) Uncovering the hidden ground state of green fluorescent protein, *Biophys. J.* 86, 168A–168A.
- Brejč, K., Sixma, T. K., Kitts, P. A., Kain, S. R., Tsien, R. Y., Ormo, M., and Remington, S. J. (1997) Structural basis for dual excitation and photoisomerization of the *Aequorea victoria* green fluorescent protein, *Proc. Natl. Acad. Sci. U.S.A.* 94, 2306–2311.
- Palm, G. J., Zdanov, A., Gaitanaris, G. A., Stauber, R., Pavlakis, G. N., and Wlodawer, A. (1997) The structural basis for spectral variations in green fluorescent protein, *Nat. Struct. Biol.* 4, 361–365.
- Stoner-Ma, D., Jaye, A. A., Matousek, P., Towrie, M., Meech, S. R., and Tonge, P. J. (2005) Observation of excited-state proton transfer in green fluorescent protein using ultrafast vibrational spectroscopy, *J. Am. Chem. Soc.* 127, 2864–2865.
- Hanson, G. T., McAnaney, T. B., Park, E. S., Rendell, M. E. P., Yarbrough, D. K., Chu, S. Y., Xi, L. X., Boxer, S. G., Montrose, M. H., and Remington, S. J. (2002) Green fluorescent protein variants as ratiometric dual emission pH sensors. 1. Structural characterization and preliminary application, *Biochemistry* 41, 15477–15488.
- McAnaney, T. B., Park, E. S., Hanson, G. T., Remington, S. J., and Boxer, S. G. (2002) Green fluorescent protein variants as ratiometric dual emission pH sensors. 2. Excited-state dynamics, *Biochemistry* 41, 15489–15494.
- McAnaney, T. B., Shi, X., Abbyad, P., Jung, H., Remington, S. J., and Boxer, S. G. (2005) Green fluorescent protein variants as ratiometric dual emission pH sensors. 3. Temperature dependence of proton transfer, *Biochemistry* 44, 8701–8711.
- Cormack, B. P., Valdivia, R. H., and Falkow, S. (1996) FACS-optimized mutants of the green fluorescent protein (GFP), *Gene* 173, 33–38.
- Elslinger, M. A., Wachter, R. M., Hanson, G. T., Kallio, K., and Remington, S. J. (1999) Structural and spectral response of green fluorescent protein variants to changes in pH, *Biochemistry* 38, 5296–5301.
- De Giorgi, F., Lartigue, L., Bauer, M. K., Schubert, A., Grimm, S., Hanson, G. T., Remington, S. J., Youle, R. J., and Ichas, F. (2002) The permeability transition pore signals apoptosis by directing Bax translocation and multimerization, *FASEB J.* 16, 607–609.
- Kissinger, C. R., Gehlhaar, D. K., and Fogel, D. B. (1999) Rapid automated molecular replacement by evolutionary search, *Acta Crystallogr. Sect. D: Biol. Crystallogr.* 55, 484–491.
- Tronrud, D. E., Teneyck, L. F., and Matthews, B. W. (1987) An efficient general-purpose least-squares refinement program for macromolecular structures, *Acta Crystallogr. Sect. A* 43, 489–501.
- Jones, T. A., Zou, J. Y., Cowan, S. W., and Kjeldgaard, M. (1991) Improved methods for building protein models in electron-density maps and the location of errors in these models, *Acta Crystallogr. Sect. A* 47, 110–119.
- Emsley, P., and Cowtan, K. (2004) Coot: model-building tools for molecular graphics, *Acta Crystallogr. Sect. D: Biol. Crystallogr.* 60, 2126–2132.
- Sheldrick, G. M., and Schneider, T. R. (1997) SHELXL: high-resolution refinement, *Macromol. Crystallogr. Part B* 277, 319–343.



24. Ormo, M., Cubitt, A. B., Kallio, K., Gross, L. A., Tsien, R. Y., and Remington, S. J. (1996) Crystal structure of the *Aequorea victoria* green fluorescent protein, *Science* 273, 1392–1395.
25. Laskowski, R. A., Macarthur, M. W., Moss, D. S., and Thornton, J. M. (1993) Procheck-a program to check the stereochemical quality of protein structures, *J. Appl. Crystallogr.* 26, 283–291.
26. van Thor, J. J., Zanetti, G., Ronayne, K. L., and Towrie, M. (2005) Structural events in the photocycle of green fluorescent protein, *J. Phys. Chem. B* 109, 16099–16108.
27. Quillin, M. L., Anstrom, D. A., Shu, X. K., O’Leary, S., Kallio, K., Chudakov, D. A., and Remington, S. J. (2005) Kindling fluorescent protein from *Anemonia sulcata*: dark-state structure at 1.38 Å resolution, *Biochemistry* 44, 5774–5787.
28. Prescott, M., Ling, M., Beddoe, T., Oakley, A. J., Dove, S., Hoegh-Guldberg, O., Devenish, R. J., and Rossjohn, J. (2003) The 2.2 Å crystal structure of a pocilloporin pigment reveals a nonplanar chromophore conformation, *Structure* 11, 275–284.
29. Shu, X., Shaner, N. C., Yarbrough, C. A., Tsien, R. Y., and Remington, S. J. (2006) Novel chromophores and buried charges control color in mFruits, *Biochemistry* 45, 9639–9647.
30. Maddalo, S. L., and Zimmer, M. (2006) The role of the protein matrix in green fluorescent protein fluorescence, *Photochem. Photobiol.* 82, 367–372.
31. Rini, M., Magnes, B. Z., Pines, E., and Nibbering, E. T. J. (2003) Real-time observation of bimodal proton transfer in acid-base pairs in water, *Science* 301, 349–352.
32. Cleland, W. W., Frey, P. A., and Gerlt, J. A. (1998) The low barrier hydrogen bond in enzymatic catalysis, *J. Biol. Chem.* 273, 25529–25532.
33. Cleland, W. W., and Kreevoy, M. M. (1994) Low-barrier hydrogen-bonds and enzymatic catalysis, *Science* 264, 1887–1890.
34. Mock, W. L., and Morsch, L. A. (2001) Low barrier hydrogen bonds within salicylate mono-anions, *Tetrahedron* 57, 2957–2964.
35. Zhao, Q. J., Abeygunawardana, C., Talalay, P., and Mildvan, A. S. (1996) NMR evidence for the participation of a low-barrier hydrogen bond in the mechanism of delta(5)-3-ketosteroid isomerase, *Proc. Natl. Acad. Sci. U.S.A.* 93, 8220–8224.
36. Hibbert, F., and Emsley, J. Hydrogen bonding and chemical reactivity, *Adv. Phys. Org. Chem.* 26, 255–380.
37. Mildvan, A. S., Harris, T. K., and Abeygunawardana, C. (1999) Nuclear magnetic resonance methods for the detection and study of low-barrier hydrogen bonds on enzymes, *Methods Enzymol.* 308, 219–245.

BI7009037